

*Citation for published version:*

Lakin, R, Barrett, PM, Stevenson, C, Thomas, RJ & Wills, M 2020, 'First evidence for a latitudinal body mass effect in extant Crocodylia and the relationships of their reproductive characters', *Biological Journal of the Linnean Society*, vol. 129, no. 4, pp. 875-887. <https://doi.org/10.1093/biolinnean/blz208>

*DOI:*

[10.1093/biolinnean/blz208](https://doi.org/10.1093/biolinnean/blz208)

*Publication date:*

2020

*Document Version*

Peer reviewed version

[Link to publication](#)

**University of Bath**

## **Alternative formats**

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 First evidence for a latitudinal body mass effect in  
2 extant Crocodylia and the relationships of their  
3 reproductive characters

4  
5  
6 Rebecca J. Lakin<sup>1, 2\*</sup>, Paul M. Barrett<sup>2</sup>, Colin Stevenson<sup>3</sup>, Robert J. Thomas<sup>4</sup>  
7 & Matthew A. Wills<sup>1</sup>

8  
9 <sup>1</sup>*Milner Centre for Evolution, University of Bath, Bath, UK BA2 7AY*, <sup>2</sup>*Department of*  
10 *Earth Sciences, Natural History Museum, Cromwell Rd, South Kensington, London SW7*  
11 *5BD*, <sup>3</sup>*Crocodyles of the World, Burford Rd, Brize Norton, Carterton OX18 3NX*,  
12 <sup>4</sup>*University of Cardiff, Cardiff, CF10 3AT*

13  
14  
15 Suggested RH – Crocodylian reproductive traits

16  
17 \*Corresponding author (r.j.lakin@bath.ac.uk)

18

19

## Abstract

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

Relationships between distribution patterns and body size have been documented in many endothermic taxa. However, the evidence for these trends in ectotherms generally is equivocal, and there have been no studies of effects in crocodylians specifically. Here, we examine the relationship between latitudinal distribution and body mass in 20 extant species of crocodylians, as well as the relationships between seven important reproductive variables. Using phylogenetically independent contrasts to inform generalised linear models, we provide the first evidence of a latitudinal effect on adult female body mass in crocodylians. In addition, we explore the relationships between reproductive variables including egg mass, hatchling mass and clutch size. We report no correlation between egg mass and clutch size, upholding previously reported within-species trends. We also find no evidence of a correlation between measures of latitudinal range and incubation temperature, contrasting with the trends found in turtles..


Keywords: *crocodylians; reproduction; generalised linear model; latitude; body size; biogeography*

## 1 Introduction

The broad impact of climate on the body size of numerous vertebrate and invertebrate groups has been demonstrated across latitudinal (Ray 1960; Wooller et al. 1985; Graves 1991; Kaspari and Vargo 1995; Saunders and Tarling 2018) and altitudinal (Bernadou et al. 2016; Davis and Burt Jr. 2019; Yu et al. 2019) gradients, insofar as these correlate with temperature gradients. While evidence of latitudinal size gradients is common in endotherms (Ashton 2002a; Blackburn and Hawkins 2004), the effect of latitude on ectotherm body sizes is more controversial (Ashton and Feldman 2003; Pincheira-Donoso et al. 2008). For example, amphibians (Ashton 2002b; Adams and Church 2008), lizards (Ashton and Feldman 2003; Cruz et al. 2005; Pincheira-Donoso et al. 2008) and turtles (Ashton and Feldman 2003; Lindeman 2008; Angielczyk et al. 2015) have all been the subjects of studies with a diversity of findings. Strikingly, however, no previous studies have attempted to test whether this correlated variation applies to another major group of living reptiles, the crocodylians.

Crocodylians are a vital component of their ecosystems, often acting as keystone species in tropical regions (Ashton 2010). They represent the last surviving pseudosuchian archosaurs, a clade that once inhabited every continent and that has persisted for at least 230 Ma (Irmis et al. 2013; Turner et al. 2017). Having survived the devastating Cretaceous-Paleogene (K-Pg) mass extinction (Bronzati et al. 2015; Puértolas-Pascual et al. 2016), as well as a number of other,

60 smaller or more protracted major extinction events (Hutchison 1982; Markwick 1993; Toljagić  
61 and Butler 2013; Mannion et al. 2015), crocodylians have demonstrated remarkable resilience to  
62 cataclysmic climate change and habitat loss (Toljagić and Butler 2013). Despite this, half of all  
63 extant crocodylian species are currently threatened with extinction and, at the current rate,  
64 vertebrate species loss will soon equal or even exceed that of the K-Pg event (Barnosky et al.  
65 2011; Pievani 2014; Lécuyer 2018). To understand possible future trends in crocodylian diversity  
66 more fully, it is important that their current diversity, ecology and distribution are fully  
67 characterized (Figure 1). Despite previous work in this area (Martin 2007; Pierce et al. 2008;  
68 Nicolaï and Matzke 2019), the nature of the relationship between geographical distribution and  
69 the reproductive biology of living crocodylians is unclear.



70  
71 *Figure 1. A representation of the relationship between the mean adult female body masses of 23*  
72 *species of crocodylian and their latitudinal midpoint. Silhouette th scaled to body mass. (a) Asia-Pacific,*  
73 *(b) The Americas, (c) Africa. Abbreviations (Asia-Pacific): Alligator sinensis, Gavialis gangeticus,*  
74 *Crocodylus palustris, Crocodylus mindorensis, Crocodylus siamensis, Tomistoma schlegelii, Crocodylus*  
75 *porosus, Crocodylus novaeguineae, Crocodylus johnstoni. (Americas): Alligator mississippiensis,*

76 Crocodylus rhombifer, Crocodylus moreletii, Crocodylus acutus, Crocodylus intermedius, Caiman  
77 crocodilus, Paleosuchus trigonatus, Melanosuchus niger, Paleosuchus palpebrosus, Caiman latirostris,  
78 Caiman yacare (*Africa*): Crocodylus cataphractus, Osteolaemus tetraspis, Crocodylus niloticus. Mesticops  
79 leptorhynchus, Crocodylus suchus and Osteolaemus osborni were excluded due to lack of reliable female  
80 mass data.

81

82         The resilience of crocodylians to historic mass extinctions is also much debated. Previous  
83 work has implicated many different factors in their survival, including diet (Sheehan and Hansen  
84 1986), aquaticism (Robertson et al. 2013), famine resistance (Robertson et al. 2013), induction  
85 of torpor at low air or water temperatures (Almandariz 1975; Brisbin Jr. et al. 1982; Sun et al.  
86 2019) and their propensity to burrow in order to take refuge from harsh environmental conditions  
87 (Thorbjarnarson 1989; Mobaraki et al. 2015). Another hypothesis concerns their unique  
88 reproductive biology (Charreau et al. 2017). Crocodylians have no sex chromosomes and hatchling  
89 sex is thought to be fully controlled by incubation temperature. Temperature dependent sex  
90 determination (TSD) systems have a threshold that yields an approximately 1:1 ratio of males  
91 and females at equilibrium (Escobedo-Galván et al 2016). Higher temperatures produce a greater  
92 proportion of males, whereas lower temperatures produce more females (Deeming and Ferguson  
93 1989; González et al. 2019). Counter-intuitively, the highest tolerable incubation temperatures  
94 produce mostly females, although these frequently fail to thrive (Marco et al. 2017). TSD was  
95 proposed as a possible buffer to extinction by Woodward and Murray (1993). Harsh  
96 environmental conditions are usually associated with a fall in ambient temperatures, and in  
97 species that utilise TSD these can result in a female-biased primary sex ratio (Tomillo et al. 2014;

2015; Carter et al. 2018). While lower temperatures may be consistent with the initial ‘nuclear winter’ effect of the end-Cretaceous bolide impact (Vellekoop et al. 2014; 2016), the longer-term warming resulting from increasing atmospheric greenhouse gas concentrations (Harrington 2001; Turner 2018) would be expected to skew sex ratios in favour of males. The latter scenario has been hypothesised as the most likely outcome of current trends in global warming (Miller et al. 2004). Although TSD has been well documented within crocodylian species (Ferguson and Joanen 1982; Deeming 2004; Piña et al. 2007), no comparisons of the effects of threshold temperature on development have been attempted across species, and little is known concerning the interrelationships between crocodylian reproductive characters.

The reproductive characters of crocodylians were first surveyed by Thorbjarnarson (1996). This work identified allometric relationships between egg mass, clutch size, clutch mass and female length: patterns that had previously been observed in other vertebrate groups (Rohwer 1988; Sinervo and Licht 1991; Shine 1992). However, the correlations and causal interactions between these reproductive variables were not the focus of this study.

Here, we used generalised linear models (GLMs) to test the relationships between the latitudinal distribution of extant crocodilian taxa and a variety of their important reproductive variables. We also tested whether there are significant relationships *between* those reproductive variables. In the context of these findings, we highlight some key questions concerning the reproductive biology, behaviour and social systems of wild crocodylians that could be addressed by future empirical and modelling work.

Figure 1: A representation of the relationship between the mean adult female body masses of 23 species of crocodylian and their latitudinal midpoint. Silhouettes scaled to body mass. Abbreviations (Asia-Pacific): *Alligator sinensis*, *Gavialis gangeticus*, *Crocodylus palustris*, *Crocodylus mindorensis*, *Crocodylus siamensis*, *Tomistoma schlegelii*, *Crocodylus porosus*, *Crocodylus novaeguineae*, *Crocodylus johnstoni*. (Americas): *Alligator mississippiensis*, *Crocodylus rhombifer*, *Crocodylus moreletii*, *Crocodylus acutus*, *Crocodylus intermedius*, *Caiman crocodilus*, *Paleosuchus trigonatus*, *Melanosuchus niger*, *Paleosuchus palpebrosus*, *Caiman latirostris*, *Caiman yacare* (Africa): *Crocodylus cataphractus*, *Osteolaemus tetraspis*, *Crocodylus niloticus*. *Masticops leptorhynchus* excluded due to lack of reliable female mass data.

## 2 Methods

### 2.1 Data Collection

We compiled data from the literature for seven morphological, environmental and reproductive variables across 24 extant crocodilian species (Appendix I). We used the species list in Grigg and Kirschner (2015) (Figure 1) and included one recently described species (*Mecistops leptorhynchus*) (Shirley 2018). Recent taxonomic uncertainty makes the attribution of some reproductive data unclear, and two species (*Crocodylus suchus* and *Osteolaemus osborni*) were



141 therefore excluded (Shirley et al. 2014; Hallmann and Griebeller 2018; Isberg et al. 2019). The  
142 data compiled were:

143

144 i. Maximum latitudinal range (°)

145

146 ii. Midpoint of latitudinal range (°)

147

148 iii. Mean mass of breeding-age female (kg)

149

150 iv. Mean fresh egg weight (g)

151

152 v. Mean hatchling mass (g)

153

154 vi. Threshold incubation temperature, or the temperature at which a clutch will produce  
155 approximately 50% males and 50% females (°C). The lower (female to male)  
156 threshold was selected in this case because females are often inviable at the higher  
157 (male to female) threshold.

158

159 vii. Mean clutch size (number of eggs laid)

160

161 viii. Mean duration of incubation (days)

162

163 We reviewed the literature in Google Scholar by searching for each species individually, coupled with  
164 appropriate terms (e.g. “*Crocodylus porosus* clutch size”, “*Caiman yacare* body mass”). Variations on  
165 these were used as necessary (e.g. “*Caiman yacare*”/ “*Caiman jacare*” + “body mass”/“female  
166 mass”/“weight”/“kg”). These were searched until pages returned only irrelevant results. Data were  
167 collated into a single spreadsheet (Appendix I). Continuous variables (female mass, egg mass,  
168 hatchling mass, clutch size and duration of incubation) were averaged to smooth over outlier results  
169 that be unrepresentative. The literature contains data from individual captive bred and wild caught  
170 specimens, as well as values that are themselves averages from various heterogeneous samples. This  
171 constrains the precision of our data, but we believe that there is a homogenous distribution of error. In  
172 addition, data on female mass were often derived from adults of different ages, and these were then  
173 averaged across multiple individuals and studies (Appendix I). Data on latitudinal range were taken  
174 from the IUCN Red List of threatened species (IUCN 2019).

175

176 For one species, *Tomistoma schlegelii*, there was insufficient information in the literature, and we  
177 therefore used data from our own observations. Hatchling mass data (see Appendix I) were  
178 averaged from a captive-born clutch laid at “Crocodiles of the World”, Oxfordshire, UK. This clutch  
179 consisted of 17 eggs, of which 14 successfully hatched (though one infant failed to thrive and died  
180 shortly after hatching). Eggs were laid on 16th April 2016 by an imported farm-raised female  
181 measuring 2.3 m (SVL). Eggs were separated and artificially incubated at 32.6° C (three eggs),  
182 32.0° C (five eggs), 31.5° C (five eggs) and 29.8° C (four eggs), and hatched after 92 days, 94

days, 95 days and 103 days, respectively. Hatchlings were weighed using a Marsden Super-SS B-100 waterproof electronic scale, accurate to 0.10 g.

## 2.2 Statistical analysis

For indicative purposes, non-phylogenetically controlled bivariate correlations between our seven variables are visualised in Figure 2. These were further tested using PGLS as sensitivity analyses. These results, and literature on archosaur reproduction generally and crocodylians specifically was used to frame five hypotheses:

- i. Adult female mass is positively correlated with latitudinal midpoint, maximum latitudinal range, egg mass, hatchling mass and/ or clutch size, as in some species of crocodylians (Verdade 2001; Campos et al. 2008; Murray et al. 2013) and birds (Wendeln 1997).
- ii. Egg mass is positively correlated with female mass, hatchling mass and/or clutch size, as in some species of crocodylians (Verdade 2001; Murray et al., 2013), birds (Wiebe and Bortolotti 1995) and turtles (Wallace et al. 2006).
- iii. Hatchling mass is positively correlated with adult female mass, latitudinal midpoint, maximum latitudinal range and/or egg mass, as in some species of crocodylians (Verdade

2001; Murray et al. 2013), birds (Ricklefs 1984; Smith et al. 1993) and turtles  
(Roosenburg and Kelley 1996; Wallace et al. 2006).

- iv. Clutch size is positively correlated with adult female mass, egg mass, latitudinal midpoint,  
maximum latitudinal range and/or duration of incubation, as in some species of  
crocodylians (Verdade 2001; Campos et al. 2008; Murray et al. 2013), birds (Erikstad et al.  
1993; Monaghan et al. 1995) and turtles (Roosenburg and Kelley 1996).
- v. Threshold incubation temperature is positively correlated with latitudinal midpoint,  
maximum latitudinal range and/or hatchling mass, as in turtles (Ewert et al. 2005).

The phylogeny of O'Brien et al. (2019) was used to produce phylogenetically-independent  
contrast (pic) values for each variable. These pic values (rather than our raw data) were used in the  
subsequent GLM analyses. This tree was constructed using multiple specimens representing  
each species and the tree was first pruned to remove duplicate specimens  
(‘Alligator\_mississippiensis2’, ‘Alligator\_mississippiensis3’, etc.). The first entry was kept by  
default in each case. Rooting and branch lengths were kept consistent with those published by  
O'Brien et al. (2019).

Before calculating the pic values, conspicuously non-normal continuous variables (female mass,  
egg mass and hatchling mass only) were log-transformed. Four species were excluded from our  
dataset because they were absent from the phylogeny of O'Brien et al. (2019) (*Crocodylus*  
*intermedius*, *Crocodylus moreletii*, *Caiman yacare* and *Masticops leptorhynchus*), leaving 20 species in  
the final analyses. We used the Akaike Information Criterion (AIC) to identify the minimum

adequate model by backward stepwise deletion of poorly fitting variables. All models used the ‘Gaussian’ error family and ‘identity’ link function, and were implemented in the statistical software environment R, using the *GGally*, *ape* and *phytools* packages (version 3.5.3) (R Core Team 2013).

We constructed five GLMs based on the phylogenetically independent contrasts of all variables. Models 3 and 5 were produced from a modified phylogeny of 17 species, excluding *Crocodylus novaeguineae*, *Osteolaemus tetraspis* and *Tomistoma schlegelii*. This was because threshold incubation temperature data were not available for these species, and also because threshold incubation temperature was shown through our Spearman’s Rank test (Figure 2) to correlate with hatchling mass (model 3), incubation duration and latitudinal values (model 5).

Model No.	Dependent variable	Independent variables
1	f.mass	Lat., egg.mass, hatch.mass, clutch.size, inc.dur
2	Egg mass	Lat, f.mass, clutch.size
3	Hatch.mass	Lat., f.mass, egg.mass, clutch.size, inc.temp, inc.dur
4	Clutch.size	Lat., f.mass, egg.mass, inc.temp.
5	Inc.temp.	Lat., egg.mass, hatch.mass, clutch.size, inc.dur

Table 1: Initial structure of GLMs analysing relationships between phylogenetically independent contrasts in the reproductive characters of extant Crocodylia. All models were conducted using the ‘gaussian’ error family and ‘identity’ link function. Abbreviations: clutch.size = mean clutch size; egg.mass = log mean fresh egg weight;

237 *f.mass* = log mean adult female body mass; *hatch.mass* = log mean hatchling mass; *inc.dur* = mean duration of  
238 incubation; *inc.temp* = threshold incubation temperature; *lat.max* = maximum latitudinal range; *t.lat.* =  
239 latitudinal midpoint.

### 3 Results

A Spearman's correlation matrix of our results is shown in Figure 2. The strongest positive correlations are between morphological reproductive characteristics (female mass vs egg mass ( $r_s = 0.85$ ,  $p < 0.0001$ ), female mass vs hatchling mass ( $r_s = 0.79$ ,  $p < 0.0001$ ), and egg mass vs hatchling mass ( $r_s = 0.94$ ,  $p < 0.0001$ )). Other correlated traits are: (1) latitudinal midpoint vs duration of incubation ( $r_s = -0.60$ ,  $p = 0.005$ ); (2) female mass vs clutch size ( $r_s = 0.67$ ,  $p = 0.002$ ); (3) egg mass vs clutch size ( $r_s = 0.56$ ,  $p = 0.01$ ); (4) hatchling mass vs clutch size ( $r_s = 0.64$ ,  $p = 0.003$ ); (5) maximum latitudinal range vs clutch size ( $r_s = 0.045$ ,  $p = 0.046$ ) (6) latitudinal midpoint vs duration of incubation ( $r_s = -0.60$ ,  $p = 0.0055$ ). PGLS analyses supported the following significant relationships: (1) female mass vs egg mass ( $p = 0.0008$ ); (2) female mass vs hatchling mass ( $p = 0.0003$ ); (3) egg mass vs hatchling mass ( $p < 0.0001$ ); (4) latitudinal midpoint vs duration of incubation ( $p = 0.007$ ); (5) female mass vs clutch size ( $p = 0.007$ ); (6) hatchling mass vs clutch size ( $p = 0.015$ ) and (7) latitudinal midpoint vs duration of incubation. Some relationships were not supported by PGLS analysis: (1) egg mass vs clutch size ( $p = 0.071$ ); (2) maximum latitudinal range vs clutch size ( $p = 0.121$ ).




Figure 2: Spearman's correlation matrix indicating positive correlations between morphological reproductive traits (female mass, egg mass, hatchling mass), and negative correlations between latitudinal midpoint, threshold incubation temperature and incubation duration. Abbreviations: clutch.size = mean clutch size; egg.mass = log mean fresh egg weight; f.mass = log mean adult female body mass; hatch.mass = log mean hatchling mass; inc.dur = mean duration of incubation; inc.temp = threshold incubation temperature; lat.max = maximum latitudinal range; t.lat. = latitudinal midpoint



### 3.1 Female Mass

Model 1 was constructed to account for the potential effects of latitudinal midpoint, maximum latitudinal range, egg mass, hatchling mass, clutch size and duration of incubation on female mass (Appendix II - initial and final models). The final model (Table 2) was the result of backward stepwise deletion based on AIC value as a measure of model efficiency. This model explained 85% of the deviance associated with adult female mass ( $D^2$ ). Mean clutch size ( $p = 0.0017$ ) was the most significant correlator to log mean female mass, followed by mean latitudinal range ( $p = 0.026$ ), log mean hatchling mass ( $0.0027$ ) and maximum latitudinal range ( $p = 0.0133$ ). Log mean egg mass and mean incubation duration both correlated poorly with log mean female mass and were dropped from the model.

Term	Coefficients	SE	F	p
t.lat	0.0421	0.0115	13.311	0.0026**
lat.max	-0.0365	0.0129	8.0132	0.0133*
log.hatch	1.1638	0.3206	13.175	0.0027**
clutch.size	0.0348	0.0090	15.047	0.0017**

*Table 2: A Generalised Linear Model (Model 1) describing the effect of latitudinal midpoint, maximum latitudinal range, hatchling mass and clutch size on the adult female mass of 20 species of crocodylian.  $D^2 = 0.8510$  SE = Standard error; F = F-value, p = p-value.*

### 3.2 Egg Mass

Model 2 was constructed to account for the potential effects of latitudinal midpoint, maximum latitudinal range, female mass and clutch size on egg mass (Appendix II). The final model (Table 3) explained 68% of the deviance associated with egg mass ( $D^2$ ). Log mean female mass ( $p < 0.0001$ ) was the only significant correlator to log mean egg mass. Latitudinal midpoint and mean clutch size correlated poorly with log mean egg mass and were dropped from the model.

Term	Coefficients	SE	F	p
max.lat	-0.0074	0.0049	1.9114	0.1858
log.f.mass	0.5117	0.0809	33.295	<0.0001***

Table 3: A Generalised Linear Model (Model 2) describing the effect of maximum latitudinal range and adult female mass on the fresh egg weight of 20 species of crocodylian.  $D^2 = 0.6776$ . SE = Standard error; F = F-value,  $p = p$ -value.

### 3.3 Hatchling Mass

Model 3 was constructed to account for the potential effects of latitudinal midpoint, maximum latitudinal range female mass, egg mass, clutch size, threshold incubation temperature and incubation duration on hatchling mass (Appendix II). The final model (Table 4) explained 90% of the deviance associated with hatchling mass ( $D^2$ ). Log mean egg mass ( $p < 0.0001$ ) was the only significant correlator to hatchling mass. Log mean female mass, latitudinal midpoint, mean clutch size and mean incubation duration correlated poorly with hatchling mass and were

dropped from the model.

Term	Coefficients	SE	F	p
lat.max	0.0052	0.0032	2.6903	0.1269
log.egg.mass	0.8348	0.0841	98.483	< 0.0001***
inc.temp	-0.0751	0.0505	2.2103	0.1629

Table 4: A Generalised Linear Model (Model 3) describing the effect of female body mass and fresh egg weight on the hatchling mass of 17 species of crocodylian.  $D^2 = 0.8997$ . SE = Standard error; F = F-value, p = p-value.

### 3.4 Clutch Size

Model 4 was constructed to account for the potential effects of latitudinal midpoint, maximum latitudinal range, female mass, egg mass and hatchling mass (Appendix II). The final model (Table 5) explained 87% of the deviance associated with clutch size ( $D^2$ ). The most significant correlators to mean clutch size were log mean female mass ( $p < 0.0001$ ) and maximum latitudinal range ( $p < 0.0001$ ), followed by latitudinal midpoint ( $p = 0.0007$ ) and mean duration of incubation ( $p = 0.107$ ). Egg mass and hatchling mass were found to correlate poorly with clutch size, and were dropped from the model.

Term	Coefficients	SE	F	p
log.f.mass	11.9056	2.0480	33.796	<0.0001 ***
t.lat	-0.8362	0.1955	18.291	0.0007***

lat.max	1.0462	0.1804	33.623	<0.0001***
inc.dur	0.2435	0.1414	2.9650	0.1071 *

Table 5: A Generalised Linear Model (Model 4) describing the effect of adult female mass, latitudinal midpoint, maximum latitudinal range and incubation duration on the clutch sizes of 20 species of crocodylian.  $D^2 = 0.8662$ . SE = Standard error; F = F-value, p = p-value.

### 3.5 Threshold Incubation Temperature

Model 5 was constructed to account for the potential effects of latitudinal midpoint, maximum latitudinal range, female mass, egg mass, hatchling mass, clutch size and incubation duration on threshold incubation temperature (Appendix II). The final model (Table 6) explained 42% of the deviance associated with threshold incubation temperature ( $D^2$ ). The only significant correlator to threshold incubation temperature was mean incubation duration ( $p = 0.0065$ ). Log mean female mass, log mean egg mass, log mean hatchling mass, mean clutch size, latitudinal midpoint and maximum latitudinal range all correlated poorly with threshold incubation temperature and were dropped from the model.


Term	Coefficients	SE	F	p
inc.dur	-0.0260	0.0081	10.201	0.0065**

Table 6: A Generalised Linear Model (Model 5) describing the effects of maximum latitudinal range, fresh egg weight, hatchling mass and duration of incubation on the threshold incubation temperature of 17 species of crocodylian.  $D^2 = 0.4215$  SE = Standard error; F = F-value, p = p-value.

## 4 Discussion

### 4.1 General overview

A robust, positive correlation between latitude and body size has been well documented in endotherms (Gillman et al. 2009; Torres-Romero et al. 2016), but is much less compelling among ectotherms (Ashton and Feldman 2003; Lindeman 2008; Cruz et al. 2005), and has never been tested for in crocodylians. Here, we demonstrate that, despite low overall correlation coefficients, both latitudinal midpoint and maximum latitudinal range correlate significantly with adult female mass once the effects of phylogeny and other potentially confounding reproductive variables are controlled for ( $p = 0.0026$  and  $0.0133$  respectively, see Model 1, section 3.1). Furthermore, we describe significant relationships between some important morphological reproductive variables across crocodylians as a whole, a phenomenon that has previously been described only within species (Verdade 2001; Campos et al. 2008; Murray et al. 2013). We demonstrate that these relationships are discrete with, for example, a significant effect of female mass on egg mass, and of egg mass on hatchling mass, but no direct effect of female mass on hatchling mass. Female mass is a significant predictor of clutch size in crocodylians, as well as both the latitudinal midpoint and maximum latitudinal range. We did not detect any significant relationship between threshold incubation temperature and either latitudinal midpoint or maximum latitudinal range (see Model 5, section 3.5). Some of the most biologically significant relationships described by our models are presented in Figure 3.



350

351 *Figure 3: Map of the interactions between some important reproductive variables in Crocodylia. Positive*  
352 *relationships shown are between latitudinal midpoint and log mean female mass ( $p = 0.0026$ ); maximum latitudinal*  
353 *range and log mean female mass ( $p = 0.013$ ); log mean egg mass and log mean female mass ( $p < 0.0001$ ); log mean*  
354 *egg mass and log mean hatchling mass ( $p < 0.0001$ ); log mean female mass and mean clutch size ( $p < 0.0001$ ;*  
355 *maximum latitudinal range and mean clutch size ( $p < 0.0001$ ); latitudinal midpoint and mean clutch size ( $p =$*   
356  *$0.0007$ ); and mean incubation duration and threshold incubation temperature ( $p = 0.0065$ ).*

357

#### 358 4.2 Implications of results

359 The strongest associations in our study are between morphological reproductive variables (Models  
360 1, 2 and 3). These include adult female mass, fresh egg weight, hatchling mass and clutch size  
361 (Figure 3). Intraspecific relationships of this kind have previously been documented in the

American alligator (*Alligator mississippiensis*) (Murray et al. 2013) and in the broad-snouted, spectacled and Yacare caimans (*Caiman latirostris*, *C. crocodilus* and *C. yacare*) (Verdade 2001; Larriera et al. 2004; Campos et al. 2008). Here, we verify the same trends across extant crocodylians as a whole. We report a significant positive effect of female mass upon clutch size (Model 4), a result that contrasts with the findings of previous species-level studies (Verdade 2001). We find no evidence for any correlation between egg mass and clutch size, which is a biological trade-off that has been described in many other taxa (Lack 1967; Sinervo and Licht 1991; Rowe 1994).

We found no significant relationship between threshold incubation temperature and hatchling mass (Models 3 and 5), suggesting that the temperature required to produce an even sex ratio is not significantly different in crocodylian species that produce large hatchlings compared with those that produce small hatchlings. This result is consistent with previous studies that examined trends within species (Joanen and McNease 1989; Campos 1993). The effects of contemporary climatic warming on the reproductive biology of crocodylians has yet to be examined (Zhang et al. 2009; Simoncini et al. 2014), but based on deep-time associations between crocodylian distributions and climate change (Markwick 1998; Quintero and Wiens 2013; Mannion et al. 2015), it is plausible that all crocodylians are vulnerable to rapid warming, and that this should be considered a threat to local populations (Dudgeon 2014).

Our results suggest that the only significant correlator to threshold incubation temperature in

crocodylians is incubation duration (Model 5). This factor may help to explain the diversity of threshold incubation temperatures as, within species, eggs incubated at higher temperatures tend to hatch sooner than those incubated at cooler temperatures (see section 2.1 on *Tomistoma schlegelii*, Methods).

We identified no significant relationships between the threshold incubation temperature and either the mean or the maximum latitudinal range of crocodylians (Model 5). This suggests that the nest temperature of crocodylians is affected by factors other than latitudinal distribution. This stands in contrast to the relationship observed in turtles, in which latitude has a significant effect on reproductive traits and nest temperature (Iverson et al. 1993; Ewert et al. 2005). The clutch size of turtles correlates with latitude (Iverson et al. 1993) and their sex ratio is affected by annual temperature fluctuations (Janzen 1994). This supports observed differences in the nest functions and parental roles of turtles and crocodylians, and suggests that the presence of adult crocodylians at the nest plays a much greater role in the temperature regulation of their nests than in turtles. Our results suggest that other factors (possibly the position, construction, location, shape or size of the nest) could contribute to the maintenance of an acceptable nest temperature (see also Grigg 1987, Brazaitis and Wantanabe 2011). In turtles, nest temperature is determined by ambient temperature and climate to a much higher degree (Hays et al. 2010; Refsnider et al. 2013; Santidrián et al. 2015). Future research on the behaviour of nesting crocodylians may be necessary to shed further light on nest temperature regulation.



### 4.3 Limitations and future work

We note that some sources in the literature measured female mass at first laying, while others measured older females. Because crocodylians grow consistently post sexual maturity (Seymour et al. 2013), this could produce variable results. While our data were means from all of the published studies we were able to identify, we recognise the limitations of literature-based data collection, and acknowledge that the standardized collection of primary data would be preferable. This use of species mean values for mass, latitudinal distribution and clutch size also masks both population level and biogeographical variation (Simoncini et al. 2009). Gathering these additional data would offer greater power, and enable both within- and cross-species comparisons.

An extension of the present study could include additional parameters such as nest dimensions, nest humidity, geographic range, and postcopulatory reproductive behaviours (nest defence, guarding of hatchlings, etc.) to test the influence of these on crocodylian distribution and behaviour.

Mortality rates among unguarded eggs and hatchlings have been suggested as factors that may promote selection for parental care (Klug and Bonsall 2010; Klug et al. 2012). Since there seems to be a wide within-species range of parenting behaviours in crocodylians (Hunt and Watanabe 1982; Platt et al. 2008), the group is a good model system for investigating the effects of different types of care. For example - do female crocodylians living in high-stress

environments (at the edge of their range or in human-dominated landscapes) exhibit the same rates and types of parenting behaviour as those living in highly protected habitats? This represents an interesting subject for future research, especially as the habitats of crocodylians are increasingly subject to human encroachment and climatic stress (Langley 2005; Amarasinghe et al. 2015; Corvera et al. 2017).

## 5 Conclusions

We present the first evidence for a latitudinal effect on the body size of extant crocodylian species (as adults and hatchlings) and make the novel observation that major reproductive characteristics of extant crocodylians follow a consistent pattern of effect across the entire order.

We report no significant relationship between either latitudinal midpoint or maximum latitudinal range and threshold incubation temperature across sampled crocodylians. This contrasts markedly with the patterns seen in turtles and raises additional questions about the parental roles of crocodylians. Further work will be needed to clarify the possible responses of crocodilian reproduction and its impact on their resilience in relation to anthropogenic warming. This may have implications for future studies on the effects of climate, latitude and life history on ectothermic amniotes, and for conservationists and government departments responsible for legislating on wildlife protection and climate change mitigation strategies.

446       6       Acknowledgements

447

448       We thank Dr. Daniel Field and Professor Tamas Szekely for their kind patience and support  
449       during the initial stages of this project. We would also like to thank Pedro L. Godroy and for his  
450       balanced and helpful commentary on our paper, and four anonymous reviewers for their kind  
451       and thorough critiques.

452

453       7       Funding Statement

454

455       This study was conducted as part of the PhD project of R. Lakin, with funding provided by the  
456       National Environmental Research Council [NE/L002434/1]. MAW thanks the Natural  
457       Environment Research Council [NE/K014951/1] and John Templeton Foundation  
458       [61408].

459

References

- Adams, D. C. and J. O. Church 2008. Amphibians do not follow Bergmann's rule. *Evolution: International Journal of Organic Evolution*, 62(2): 413–420.
- Almandarz, E.1975. The use of chilled water to transfer adult crocodilians. *International Zoo Yearbook*, 15(1): 171–172.
- Amarasinghe, A.T., Madawala, M.B., Karunarathna, D.S., Manolis, S.C., de Silva, A. and Sommerlad, R., 2015. Human-crocodile conflict and conservation implications of saltwater crocodiles *Crocodylus porosus* (Reptilia: Crocodylia: Crocodylidae) in Sri Lanka. *Journal of Threatened Taxa*, 7(5), pp.7111-7130.
- Angielczyk, K. D., R. W. Burroughs, and C. R. Feldman 2015. Do turtles follow the rules? Latitudinal gradients in species richness, body size, and geographic range area of the world's turtles. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, 324(3): 270–294.
- Ashton, K. G. 2002a. Patterns of within-species body size variation of birds: strong evidence for Bergmann's rule. *Global Ecology and Biogeography*, 11(6): 505–523.
- Ashton, K. G. 2002b. Do amphibians follow Bergmann's Rule? *Canadian Journal of Zoology*, 80(4): 708-716

481 Ashton, K. G. and C. R. Feldman 2003. Bergmann's rule in nonavian reptiles: turtles follow it,  
 482 lizards and snakes reverse it. *Evolution*, 57(5): 1151–1163.

483

484 Ashton, P. J. 2010. The demise of the Nile crocodile (*Crocodylus niloticus*) as a keystone species for  
 485 aquatic ecosystem conservation in South Africa: The case of the Olifants River. *Aquatic*  
 486 *Conservation: Marine and Freshwater Ecosystems* 20(5): 489–493.

487

488 Barnosky, A. D., N. Matzke, S. Tomiya, G. O. U. Wogan, B. Swartz, T. B. Quental, C. Marshall,  
 489 J. L. McGuire, E. L. Lindsey, K. C. Maguire, B. Mersey, and E. A. Ferrer 2011. Has the  
 490 Earth's sixth mass extinction already arrived? *Nature*, 471(7336): 51.

491

492 Bernadou, A., C. Roemermann, N. Gratiashvili, and J. Heinze 2016. Body size but not colony  
 493 size increases with altitude in the holarctic ant, *Leptothorax acervorum*. *Ecological*  
 494 *entomology*, 41(6): 733–736.

495 Blackburn, T. M. and B. A. Hawkins 2004. Bergmann's rule and the mammal fauna of northern  
 496 North America. *Ecography*, 27(6): 715–724.

497 Brazaitis, P. and M. E. Wantanabe 2011. Crocodilian behaviour: a window to dinosaur  
 498 behaviour?. *Historical Biology*, 23(1): 73–90.

499

500 Brisbin Jr., I. L., E. A. Standora, and M. J. Vargo 1982. Body temperatures and behavior of

501 American alligators during cold winter weather. *American Midland Naturalist*, P. 209–  
502 218.

503

504 Bronzati, M., F. C. Montefeltro, and M. C. Langer 2015. Diversification events and the effects of  
505 mass extinctions on Crocodyliformes evolutionary history. *Royal Society open science*,  
506 2(5): 140385.

507 Campos, Z. 1993. Effect of habitat on survival of eggs and sex ratio of hatchlings of *Caiman*  
508 *crocodilus yacare* in the Pantanal, Brazil. *Journal of Herpetology*, 127–132.

509 Campos, Z., Llobet, A.Q., Piña, C.I. and Magnusson, W.E., 2010. Yacare caiman (*Caiman*  
510 *yacare*). *Crocodyles. Status survey and conservation action plan*, 3; 23-28.

511 Campos, Z., W. Magnusson, T. Sanaiotti, and M. Coutinho 2008. Reproductive trade-offs in *Caiman*  
512 *crocodilus crocodilus* and *Caiman crocodilus yacare*: implications for size-related management  
513 quotas. *The Herpetological Journal*, 18(2): 91–96.

514 Carter, A.W., Sadd, B.M., Tuberville, T.D., Paitz, R.T. and Bowden, R.M., 2018. Short  
515 heatwaves during fluctuating incubation regimes produce females under temperature-  
516 dependent sex determination with implications for sex ratios in nature. *Scientific*  
517 *Reports*, 8(1); 3.

518 Charruau, P., Cantón, D.A.M.G. and Mendez de la Cruz, F.R., 2017. Additional details on  
519 temperature-dependent sex determination in *Crocodylus acutus*. *Salamandra* 53(2); 304-  
520 308. Cleary, T. J., R. B. Benson, S. E. Evans, and P. M. Barrett 2018. Lepidosaurian  
521 diversity in the Mesozoic–Palaeogene: the potential roles of sampling biases and

522 environmental drivers. *Royal Society open science*, 5(3): 171830.

523 Corvera, M.D., Manalo, R.I. and Aquino, M.T.R., 2017. People and Crocodiles Sharing One  
 524 Environment: An Analysis of Local Human-Crocodile Conflict Management Strategies  
 525 in the Philippines. *Journal of Animal Science and Research*, 1(1): 1-6.

526

527 Cruz, F. B., L. A. Fitzgerald, R. E. Espinoza, and J. A. Schulte Ii 2005. The importance of  
 528 phylogenetic scale in tests of Bergmann's and Rapoport's rules: lessons from a clade of  
 529 South American lizards. *Journal of Evolutionary Biology*, 18(6): 1559–1574.

530

531 Davis, K. M. and E. H. Burt Jr 2019. Size increase with altitude in the Rufous-collared Sparrow  
 532 (*Zonotrichia capensis*). *The Wilson Journal of Ornithology*. 131(1): 135-139

533 Deeming, D. C. 2004. Prevalence of TSD in crocodilians. *Temperature-dependent sex determination in*  
 534 *vertebrates*, 33–41.

535 Deeming, D. C. and M. W. J. Ferguson 1989. The mechanism of temperature dependent sex  
 536 determination in crocodilians: a hypothesis. *American Zoologist*, 29(3): 973–985.

537 Dudgeon, D., 2014. Threats to freshwater biodiversity in a changing world. *Global*  
 538 *Environmental Change*, 243-253.

539 Erikstad, K. E., J. O. Bustnes, and T. Moum 1993. Clutch-size determination in precocial birds: a  
 540 study of the common eider. *The Auk*, 110(3): 623–628.

541 Escobedo-Galván, A.H., López-Luna, M.A. and Cupul-Magaña, F.G., 2016. Thermal fluctuation  
542 within nests and predicted sex ratio of Morelet's Crocodile. *Journal of Thermal Biology*,  
543 58; 23-28.

544 Ewert, M. A., J. W. Lang, and C. E. Nelson 2005. Geographic variation in the pattern of  
545 temperature-dependent sex determination in the American snapping turtle (*Chelydra*  
546 *serpentina*). *Journal of Zoology*, 265(1): 81–95.

547 Ferguson, M. W. J. and T. Joanen 1982. Temperature of egg incubation determines sex in *Alligator*  
548 *mississippiensis*. *Nature*, 296(5860): 850.

549 Gillman, L. N., D. J. Keeling, H. A. Ross, and S. D. Wright 2009. Latitude, elevation and the  
550 tempo of molecular evolution in mammals. *Proceedings of the Royal Society B: Biological*  
551 *Sciences*, 276(1671): 3353–3359.

552 González et al. 2019. The sex-determination pattern in crocodilians: a systematic review of three  
553 decades of research. *Journal of Animal Ecology* 88(9): 11417-1427.

554 Graves, G. R. 1991. Bergmann's rule near the equator: Latitudinal clines in body size of an Andean  
555 passerine bird. *Proceedings of the National Academy of Sciences*, 88(6): 2322–2325.

556 Grigg, G.C. 1987. Water relations of crocodilian eggs: management considerations. IN:  
557 Grahame, J. W. W., Manolis, S. C. and Whitehead, P. (eds.) *Wildlife Management:*  
558 *Crocodiles and Alligators*, Chipping Norton: S. Beatty and Sons Pty Ltd, 499-502.

559 Hallmann, K. and Griebeler, E.M., 2018. An exploration of differences in the scaling of life



560 history traits with body mass within reptiles and between amniotes. *Ecology and*  
561 *Evolution*, 8(11); 5480-5494.

562 Harrington, G.J., 2001. Impact of Paleocene/Eocene greenhouse warming on North American  
563 paratropical forests. *Palaaios*, 16(3), 266-278.

564 Hays, G. C., S. Fossette, K. A. Katselidis, G. Schofield, and M. B. Gravenor 2010. Breeding  
565 periodicity for male sea turtles, operational sex ratios, and implications in the face of  
566 climate change. *Conservation Biology*, 24(6): 1636–1643.

567 Hunt, R. H. and M. E. Watanabe 1982. Observations on maternal behavior of the American  
568 alligator, *Alligator mississippiensis*. *Journal of Herpetology*, 16(3): 235–239.

569 Hutchison, J. H. 1982. Turtle, crocodilian, and champsosaur diversity changes in the Cenozoic of the  
570 north-central region of western United States. *Palaeogeography, Palaeoclimatology,*  
571 *Palaeoecology*, 37(2-4): 149–164.

572 Irmis, R. B., S. J. Nesbitt, and H.-D. Sues 2013. Early Crocodylomorpha. *Geological Society,*  
573 *London, Special Publications*, 379(1): 275–302.

574 Isberg, S., Combrink, X., Lippai, C. and Balaguera-Reina, S.A. 2019. *Crocodylus niloticus*. *The*  
575 *IUCN Red List of Threatened Species 2019*; e.T45433088A3010181.Iverson, J. B., C. P.  
576 Balgooyen, K. K. Byrd, and K. K. Lyddan 1993. Latitudinal variation in egg and clutch  
577 size in turtles. *Canadian Journal of Zoology*, 71(12): 2448–2461.

578 IUCN 2019. The IUCN Red List of Threatened Species. Version 2019-2. Accessed 04/11/18 at:

579           <https://www.iucnredlist.org>.

580   Janzen, F. J. 1994. Climate change and temperature-dependent sex determination in reptiles.

581           *Proceedings of the National Academy of Sciences*, 91(16): 7487–7490.

582

583   Joanen, T. and L. L. McNease 1989. Ecology and physiology of nesting and early development

584           of the American alligator. *American Zoologist*, 29(3): 987–998.

585

586   Kaspari, M. and E. L. Vargo 1995. Colony size as a buffer against seasonality: Bergmann’s rule in

587           social insects. *The American Naturalist*, 145(4): 610–632.

588

589   Klug, H., S. H. Alonzo, and M. B. Bonsall   2012. Theoretical foundations of parental care. In: *The*

590           *Evolution of Parental Care*, 21–36.

591

592   Klug, H. and M. B. Bonsall   2010. Life history and the evolution of parental care. *Evolution:*

593           *International Journal of Organic Evolution*, 64(3): 823–835.

594

595   Lack, D. 1967. The significance of clutch-size in waterfowl. *Wildfowl*, 18(18): 125–128.

596

597   Larriera, A., C. I. Piña, P. Siroski, and L. M. Verdade 2004. Allometry of reproduction in wild

598           broad-snouted caimans (*Caiman latirostris*). *Journal of Herpetology*, 38(2): 301–305.

599 Langley, R.L., 2005. Alligator attacks on humans in the United States. *Wilderness &*  
600 *Environmental Medicine*, 16(3), pp.119-124.

601

602 Lécuyer, C. 2018. Learning from past climatic changes. *Science*, 360(6396): 1400–1401.

603

604 Lindeman, P. V. 2008. Evolution of body size in the map turtles and sawbacks (Emydidae:  
605 Deirochelyinae: Graptemys). *Herpetologica*, 64(1): 32–46.

606 Mannion, P.D., Benson, R.B., Carrano, M.T., Tennant, J.P., Judd, J. and Butler, R.J., 2015.  
607 Climate constrains the evolutionary history and biodiversity of crocodylians. *Nature*  
608 *Communications*; 6, 8438.

609 Marco, P., Virginia, M., Leiva, P.M.D.L., Iungman, J.L., Simoncini, M.S. and Piña, C.I., 2017.  
610 New evidence characterizing Temperature-dependent Sex Determination in Broad-  
611 snouted caiman, *Caiman latirostris*. *Herpetological Conservation and Biology*, 12; 78-  
612 84. Markwick, P. J. 1993. Crocodilian diversity and distributional responses to climate  
613 changes over the last 100 Ma. *Geological Society of America, Abstracts with*  
614 *Programs;(United States)*, 25(3).

615 Markwick, P.J., 1998. Fossil crocodilians as indicators of Late Cretaceous and Cenozoic  
616 climates: implications for using palaeontological data in reconstructing palaeoclimate.  
617 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 137(3-4): 205-271.

618 Markwick, P. J. 2002. Integrating the present and past records of climate, biodiversity and  
619 biogeography: implications for palaeoecology and palaeoclimatology. *Geological Society,*

620           *London, Special Publications* 194(1): 179–199.

621  
622   Martin, S. 2007. Global diversity of crocodiles (Crocodilia, Reptilia) in freshwater. *Fresh-water*  
623           *Animal Diversity Assessment*, 587–591. Springer.

624  
625   Miller, D., J. Summers, and S. Silber 2004. Environmental versus genetic sex determination: a  
626           possible factor in dinosaur extinction? *Fertility and Sterility*, 81(4): 954–964.

627  
628   Mobaraki, A., M. R. Silva, and E. Abtin 2015. Sustainable management and conservation of  
629           the mugger crocodile (*Crocodylus palustris*) in Iran. Master's thesis, International  
630           University of Andalusia.

631  
632   Monaghan,, M. Bolton, and D. Houston 1995. Egg production constraints and the evolution of avian  
633           clutch size. *Proceedings of the Royal Society of London. Series B: Biological Sciences*,  
634           259(1355): 189–191.

635  
636   Murray, C. M., M. Easter, M. Merchant, A. Cooper, and B. I. Crother 2013. Can reproductive  
637           allometry assess population marginality in crocodilians? A comparative analysis of Gulf  
638           Coast American Alligator (*Alligator mississippiensis*) populations. *Copeia* 2013(2): 268–  
639           276.

- 641 Nicholson, D. B., P. A. Holroyd, P. Valdes, and P. M. Barrett 2016. Latitudinal diversity gradients  
642 in Mesozoic non-marine turtles. *Royal Society open science*, 3(11): 160581.
- 643 Nicolai, M.P. and Matzke, N.J. (2019) Trait-based range expansion aided in the global radiation  
644 of Crocodylidae. *Global Ecology and Biogeography*, in print.
- 645
- 646 O'Brien, H. D., L. M. Lynch, K. A. Vliet, J. Brueggen, G. M. Erickson, and P. M. Gignac 2019.  
647 Crocodylian head width allometry and phylogenetic prediction of body size in extinct  
648 crocodyliforms. *Integrative Organismal Biology*, in press.
- 649 Pierce, S. E., K. D. Angielczyk, and E. J. Rayfield 2008. Patterns of morphospace occupation and  
650 mechanical performance in extant crocodilian skulls: a combined geometric  
651 morphometric and finite element modeling approach. *Journal of Morphology*, 269(7):  
652 840–864.
- 653
- 654 Pievani, T. 2014. The sixth mass extinction: Anthropocene and the human impact on biodiversity.  
655 *Rendiconti Lincei*, 25(1): 85–93.
- 656
- 657 Piña, C., A. Larriera, P. Siroski, L. Verdade, and V. Lance 2007. The temperature-sensitive period  
658 (TSP) during incubation of broad-snouted caiman (*Caiman latirostris*) eggs. *Amphibia-*  
659 *Reptilia*, 28(1): 123–128.
- 660

661 Pincheira-Donoso, D., D. J. Hodgson, and T. Tregenza 2008. The evolution of body size under  
 662 environmental gradients in ectotherms: why should Bergmann's rule apply to lizards?  
 663 *BMC Evolutionary Biology*, 8(1): 68.  
 664

665 Platt, S. G., T. R. Rainwater, J. B. Thorbjarnarson, and S. T. McMurry 2008. Reproductive  
 666 dynamics of a tropical freshwater crocodilian: Morelet's crocodile in northern Belize.  
 667 *Journal of Zoology*, 275(2): 177–189.  
 668

669 Puértolas-Pascual, E., A. Blanco, C. A. Brochu, and J. I. Canudo 2016. Review of the Late  
 670 Cretaceous-early Paleogene crocodylomorphs of Europe: Extinction patterns across the K-Pg  
 671 boundary. *Cretaceous Research*, 57: 565–590.

672 Quintero, I. and Wiens, J. J., 2013. Rates of projected climate change dramatically exceed past  
 673 rates of climatic niche evolution among vertebrate species. *Ecology Letters*, 16(8): 1095-  
 674 1103.  
 675

676 R Core Team 2013. *R: A Language and Environment for Statistical Computing*. R Foundation for  
 677 Statistical Computing, Vienna, Austria.  
 678

679 Ray, C. 1960. The application of Bergmann's and Allen's rules to the poikilotherms. *Journal*  
 680 *of morphology*, 106(1): 85–108.  
 681

- 682 Refsnider, J. M., B. L. Bodensteiner, J. L. Reneker, and F. J. Janzen 2013. Nest depth may not  
683 compensate for sex ratio skews caused by climate change in turtles. *Animal Conservation*,  
684 16(5): 481–490.
- 685 Ricklefs, R. E. 1984. Egg dimensions and neonatal mass of shorebirds. *The Condor*, 86(1): 07–11.
- 686 Robertson, D. S., W. M. Lewis, P. M. Sheehan, and O. B. Toon 2013. K-Pg extinction patterns in  
687 marine and freshwater environments: The impact winter model. *Journal of Geophysical*  
688 *Research: Biogeosciences*, 118(3): 1006–1014.
- 689 Rohwer, F. C. 1988. Inter and intraspecific relationships between egg size and clutch size in  
690 waterfowl. *The Auk*, 105(1): 161–176.
- 691 Roosenburg, W. M. and K. C. Kelley 1996. The effect of egg size and incubation temperature on  
692 growth in the turtle, *Malaclemys terrapin*. *Journal of Herpetology*, 30(2): 198–204.
- 693 Rowe, J. W. 1994. Reproductive variation and the egg size-clutch size trade-off within and  
694 among populations of painted turtles (*Chrysemys picta bellii*). *Oecologia*, 99(1-2): 35–44.
- 695 Santidrián T. P., M. Genovart, F. V. Paladino, J. R. Spotila, and D. Oro 2015. Climate change  
696 overruns resilience conferred by temperature-dependent sex determination in sea turtles and  
697 threatens their survival. *Global Change Biology*, 21(8): 2980–2988.
- 698 Saunders, R. A. and G. A. Tarling 2018. Southern Ocean mesopelagic fish comply with  
699 Bergmann’s rule. *The American Naturalist* 191(3): 343–351.
- 700 Seymour, R. S., C. Gienger, M. L. Brien, C. R. Tracy, S. C. Manolis, G. J. Webb, and K. A.

701 Christian 2013. Scaling of standard metabolic rate in estuarine crocodiles *Crocodylus*  
702 *porosus*. *Journal of Comparative Physiology B*, 183(4): 491–500.

703 Sheehan, P. M. and T. A. Hansen 1986. Detritus feeding as a buffer to extinction at the end of  
704 the Cretaceous. *Geology*, 14(10): 868–870.

705 Shine, R. 1992. Relative clutch mass and body shape in lizards and snakes: is reproductive  
706 investment constrained or optimized? *Evolution*, 46(3): 828–833.

707 Shirley, M.H., Carr, A.N., Nestler, J.H., Vliet, K.A. and Brochu, C.A., 2018. Systematic revision  
708 of the living African slender-snouted crocodiles (*Mecistops* Gray, 1844). *Zootaxa*,  
709 4504(2); 151-193.

710 Simoncini, M. S., F. B. Cruz, A. Larriera, and C. I. Piña 2014. Effects of climatic conditions on  
711 sex ratios in nests of broad-snouted caiman. *Journal of Zoology*, 293(4): 243–251.

712 Simoncini, M. S., C. I. Piña, and P. A. Siroski 2009. Clutch size of *Caiman latirostris*  
713 (Crocodylia: Alligatoridae) varies on a latitudinal gradient. *North-Western Journal of*  
714 *Zoology*, 5(1): 191.

715 Sinervo, B. and P. Licht 1991. Proximate constraints on the evolution of egg size, number, and  
716 total clutch mass in lizards. *Science*, 252(5010): 1300–1302.

717 Smith, H. G., U. Ottosson, and T. Ohlsson 1993. Interclutch variation in egg mass among starlings  
718 *Sturnus vulgaris* reflects female condition. *Ornis Scandinavica*, 311–316.

719 Sun, H., X. Zuo, L. Sun, P. Yan, F. Zhang, H. Xue, E. Li, Y. Zhou, R. Wu, and X. Wu 2019.



720           Insights into the seasonal adaptive mechanisms of Chinese alligators (*Alligator sinensis*)  
721           from transcriptomic analyses. *Australian Journal of Zoology*, 66(2): 93–102.

722   Thorbjarnarson, J. B. 1989. Ecology of the American crocodile (*Crocodylus acutus*). *Crocodiles:*  
723           *Their ecology, management, and conservation*, 228–258.

724   Thorbjarnarson, J. B. 1996. Reproductive characteristics of the order Crocodylia. *Herpetologica*, P.  
725           8–24.

726   Toljagić, O. and R. J. Butler 2013. Triassic–Jurassic mass extinction as trigger for the Mesozoic  
727           radiation of crocodylomorphs. *Biology Letters*, 9(3): 20130095.

728   Tomillo, P.S., Genovart, M., Paladino, F.V., Spotila, J.R. and Oro, D., 2015. Climate change  
729           overruns resilience conferred by temperature-dependent sex determination in sea turtles  
730           and threatens their survival. *Global Change Biology*, 21(8), pp.2980-2988.

731   Tomillo, P.S., Oro, D., Paladino, F.V., Piedra, R., Sieg, A.E. and Spotila, J.R., 2014. High beach  
732           temperatures increased female-biased primary sex ratios but reduced output of female  
733           hatchlings in the leatherback turtle. *Biological Conservation*, 176; 71-79. Torres-Romero,  
734           E. J., I. Morales-Castilla, and M. Á. Olalla-Tárraga 2016. Bergmann’s rule in the oceans?  
735           Temperature strongly correlates with global interspecific patterns of body size in marine  
736           mammals. *Global Ecology and Biogeography*, 25(10): 1206–1215.

737   Turner, S.K., 2018. Constraints on the onset duration of the Paleocene–Eocene Thermal  
738           Maximum. *Philosophical Transactions of the Royal Society A: Mathematical, Physical*  
739           *and Engineering Sciences*, 376(2130), p.20170082.

- 740 Turner, A.H., Pritchard, A.C. and Matzke, N.J., 2017. Empirical and Bayesian approaches to  
741 fossil-only divergence times: a study across three reptile clades. *PloS ONE*, 12(2);  
742 e0169885.
- 743 Vellekoop, J., Esmeray-Senlet, S., Miller, K.G., Browning, J.V., Sluijs, A., van de Schootbrugge,  
744 B., Damsté, J.S.S. and Brinkhuis, H., 2016. Evidence for Cretaceous-Paleogene  
745 boundary bolide “impact winter” conditions from New Jersey, USA. *Geology*, 44(8),  
746 pp.619-622.
- 747 Vellekoop, J., Sluijs, A., Smit, J., Schouten, S., Weijers, J.W., Damsté, J.S.S. and Brinkhuis, H.,  
748 2014. Rapid short-term cooling following the Chicxulub impact at the Cretaceous–  
749 Paleogene boundary. *Proceedings of the National Academy of Sciences*, 111(21),  
750 pp.7537-7541.
- 751 Verdade, L. M. 2001. Allometry of reproduction in broad-snouted caiman (*Caiman latirostris*).  
752 *Brazilian Journal of Biology*, 61(3): 431–435.
- 753 Wallace, B. P., P. R. Sotherland, P. S. Tomillo, S. S. Bouchard, R. D. Reina, J. R. Spotila, and F.  
754 V. Paladino 2006. Egg components, egg size, and hatchling size in leatherback turtles.  
755 *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 145(4):  
756 524–532.
- 757
- 758 Wendeln, H. 1997. Body mass of female common terns (*Sterna hirundo*) during courtship:  
759 relationships to male quality, egg mass, diet, laying date and age. *Colonial Waterbirds*, P.

760 235–243.

761

762 Wiebe, K. and G. Bortolotti 1995. Egg size and clutch size in the reproductive investment of

763 American kestrels. *Journal of Zoology*, 237(2): 285–301.

764

765 Woodward, D. E. and J. D. Murray 1993. On the effect of temperature-dependent sex determination

766 on sex ratio and survivorship in crocodilians. *Proceedings of the Royal Society of London.*

767 *Series B: Biological Sciences*, 252(1334): 149–155.

768

769 Wooller, R. D., D. A. Saunders, J. S. Bradley, and C. P. de Rebeira 1985. Geographical variation

770 in size of an Australian honeyeater (Aves: Meliphagidae): an example of Bergmann's

771 rule. *Biological Journal of the Linnean Society*, 25(4): 355–363.

772

773 Yu, T. L., D. L. Wang, M. Busam, and Y. H. Deng 2019. Altitudinal variation in body size in *Bufo*

774 *minshanicus* supports Bergmann's Rule. *Evolutionary Ecology*, 1–12.

775

776 Zhang, F., Y. Li, Z. Guo, and B. R. Murray 2009. Climate warming and reproduction in Chinese

777 alligators. *Animal Conservation*, 12(2): 128–137.